

Observation of candidate exotic mesons containing heavy quarks with Belle

S.-K. Choi (For Belle collaboration)
Gyeongsang Nat. University, Jinju, Rep. of Korea

We report the observation of two resonance-like structures in the $\pi^+\chi_{c1}$ invariant mass distribution near 4.1 GeV in exclusive $B \rightarrow K\pi^+\chi_{c1}$ decays. A detailed Dalitz-plot analysis demonstrates that these structures cannot be produced by reflections from any known and possibly unknown resonances in the $K\pi$ channel. If these two peaks are produced by resonances in the $\pi^+\chi_{c1}$ channel, their minimal quark structure would have to be a $c\bar{c}u\bar{d}$ tetraquark arrangement, similar to that proposed for the $Z^+(4430)$ structure reported by Belle last year in the $\pi^+\psi'$ mass distribution produced in $B \rightarrow K\pi^+\psi'$ decays. In addition, we report some new measurements on the properties of the $X(3872)$ meson and the $1^{--}Y$ states that are produced with initial state radiation. The analyses are based on a large data sample recorded at the $\Upsilon(4S)$ resonances and nearby continuum with the Belle detector at the KEKB asymmetric-energy e^+e^- collider.

1. Introduction

The B -factories were built to study CP violation in B -meson decays. They also serve as a plentiful source of $c\bar{c}$ – like quark pairs produced via the color-suppressed, but CKM-favored $b \rightarrow c\bar{c}s$ decay mode and by s -channel $e^+e^- \rightarrow c\bar{c}$ annihilation via initial state radiation. A number of particles are observed, the so-called XYZ mesons, that do not fit into any of the predicted but as yet unobserved charmonium states. As a result, they are considered possible candidates for new, exotic types of particles such as multiquark states or $c\bar{c}$ -gluon hybrids. The former include either mesonic molecules ($c\bar{q}q\bar{q}$) or diquark-diantiquark ($cq\bar{c}\bar{q}$) tetraquark mesons. We report on the recently observed Z family of states that have non-zero charge, which means they can neither be $c\bar{c}$ nor hybrid states. In addition new measurements related to the $X(3872)$ are mentioned, and properties of the $1^{--}Y$ family of states that are produced via the initial state radiation process, including those of the possible $s\bar{s}$ analog, the $Y_s(2175)$, are reported.

2. Charged resonance-like Z_1 and Z_2 states

In 2007 Belle reported the first charged charmonium-like state, denoted by $Z^+(4430)$, in the $\pi^+\psi'$ mass distribution produced in $B \rightarrow K\pi^+\psi'$ decays [1]. After vetoing the peak regions of the $K^*(890)$ and $K_2^*(1430)$ resonances in order to minimize their effects (the K^* veto), the $\pi^+\psi'$ mass distribution (shown in Fig. 1) exhibits a strong enhancement near 4.43 GeV/ c^2 . A fit with an S -wave Breit Wigner function for signal peak and a phase-spacelike function for the non-peaking background gives a mass and width of $M = 4433 \pm 4 \pm 2$ MeV/ c^2 and $\Gamma = 45_{-13}^{+18}(\text{stat})_{-13}^{+30}(\text{syst})$ MeV/ c^2 and a signal significance of 6.5σ . The product branching fraction is measured to be $B(B^0 \rightarrow K^- Z^+(4430)) \times B(Z^+(4430) \rightarrow \pi^+\psi') = (4.1 \pm 1.0 \pm 1.4) \times 10^{-5}$.

This $Z(4430)^+$ observation motivated a search for other charged charmonium-like states in $\bar{B}^0 \rightarrow K^-\pi^+c\bar{c}$ decays [2].

The left panel of Fig. 2 shows a Dalitz plot of $M^2(K^+\pi^-)$ (horizontal) versus $M^2(\pi^+\chi_{c1})$ (vertical) for $\bar{B}^0 \rightarrow K^-\pi^+\chi_{c1}$ events. Some clustering is evident in vertical bands at the mass of the $K^*(890)$ and $K_2^*(1430)$, which correspond to $B \rightarrow K^*\chi_{c1}$ decays. In addition, a clustering of events in a broad horizontal band is also apparent. To characterize this signal clustering in the Dalitz plot, Belle did a likelihood fit with a two-variable ($M^2(K^-\pi^+)$ & $M^2(\pi^+\chi_{c1})$) function that is a superposition of various interfering Breit-Wigner amplitudes that includes all known $K^-\pi^+$ resonances plus $\pi^+\chi_{c1}$ resonances. The binned fit to the Dalitz plot is performed using a very small bin size.




Figure 1: The $M(\pi^+\psi')$ distribution produced in $B \rightarrow K\pi^\pm\psi'$ decays. The K^* veto was applied. The shaded histogram shows the combinatoric background level inferred from side-band data.




Figure 2: a) The $M^2(K^-\pi^+)$ (horizontal) vs $M^2(\pi^+\chi_{c1})$ Dalitz plot distribution for the $B \rightarrow K\pi^+\chi_{c1}$ candidate events. b) The $M(\pi^+\chi_{c1})$ distribution projected from the second vertical slice in the range of $1.0 \text{ GeV}^2/c^4 < M^2(K^-\pi^+) < 1.75 \text{ GeV}^2/c^4$ in the Dalitz plot a), which is the most sensitive to Z_1 and Z_2 . The dots with error bars represent data, the solid (dashed) histogram is the Dalitz plot fit model with all known K^* and two (without any) $\pi^+\chi_{c1}$ resonances.

To display the fit results, the Dalitz plot is divided into the four vertical slices shown in Fig. 2 a) and three horizontal slices. The second vertical slice in Fig. 2 a) is the most sensitive to possible Z^+ signals; the first vertical slice has some Z^+ sensitivity. The right plot in Fig. 2 b) shows the projected $\pi^+\chi_{c1}$ mass distribution from the second vertical $K\pi$ mass slice, the region with most discrimination; the histograms are the results of fits to the entire Dalitz plot that are described below.

The dashed histogram shows the results of a fit to the $\pi^+\chi_{c1}$ structure with a fit function that includes all known $K\pi$ resonances, but no $\pi^+\chi_{c1}$ resonance. The fit CL is very poor, only 4×10^{-12} . When we include a non-resonant $K\pi\chi_{c1}$ term and an additional K^* resonance with floating mass and width into the fit function, the fit CL (from a toy MC study) is still only 2×10^{-4} . If we add one spin=0 $\pi\chi_{c1}$ resonance, the fit gives a mass and width of $M = 4150^{+31}_{-10} \text{ MeV}/c^2$ and $\Gamma = 352^{+99}_{-43} \text{ MeV}/c^2$ with signal significance of 10.7σ . However, the fit CL is only 0.1%.

If we instead use two $\pi\chi_{c1}$ resonances, denoted by Z_1 and Z_2 , the fit gives masses and widths of $M_{Z_1} = 4051 \pm 14^{+20}_{-41} \text{ MeV}/c^2$, $\Gamma_{Z_1} = 82^{+21}_{-17}(\text{stat})^{+47}_{-22}(\text{syst}) \text{ MeV}/c^2$, $M_{Z_2} = 4248^{+44}_{-29}(\text{stat})^{+180}_{-35}(\text{syst}) \text{ MeV}/c^2$ and $\Gamma_{Z_2} = 177^{+54}_{-39}(\text{stat})^{+316}_{-61}(\text{syst}) \text{ MeV}/c^2$, and a fit CL of 40%. The corresponding product branching fractions are $B(\bar{B}^0 \rightarrow$




Figure 3: a) The $M(J/\psi\pi^+\pi^-)$ distribution for the $X(3872)$ region produced in $B^0 \rightarrow J/\psi\pi^+\pi^-K^+\pi^-$ decays. b) The $K\pi$ mass distribution for $B \rightarrow X(3872)K\pi$ candidate events. The dash-dotted curve shows the non-resonant $K\pi$ component, the dotted curve the $K^*(892)^0$ component and the dashed curve the background from sideband.

$K^-Z_1^+ \times B(Z_1^+ \rightarrow \pi^+\chi_{c1}) = (3.1_{-0.8}^{+1.5}(\text{stat})_{-1.6}^{+3.7}(\text{syst})) \times 10^{-5}$ and $B(\bar{B}^0 \rightarrow K^-Z_2^+) \times B(Z_2^+ \rightarrow \pi^+\chi_{c1}) = (4.0_{-0.9}^{+2.3}(\text{stat})_{-0.5}^{+19.7}(\text{syst})) \times 10^{-5}$. These are comparable to those of other charmonium-like states as well as the $Z(4430)^+$. The two resonances structure is favored over the one resonance assumption at the 5.7σ level.

We tried to fit with many different $K\pi$ resonance options including the addition of new resonances with floating masses and widths. However, no model that only has dynamics in the $K\pi$ channel can fit the Dalitz plot.

3. News on the $X(3872)$

The threshold mass of the $D^0\bar{D}^{*0}$ system is $3871.8 \pm 0.4 \text{ MeV}/c^2$, while the PDG 2008 world average mass of the $X(3872)$ is $3871.9 \pm 0.5 \text{ MeV}/c^2$. This similarity in mass is a frequently cited characteristic of the $X(3872)$ and highly suggestive of a $D\bar{D}^*$ molecule-like structure. This motivated searches for $X(3872) \rightarrow D\bar{D}^*$ decays, and peaks with masses that are slightly higher than $3872 \text{ MeV}/c^2$ were reported both by Belle ($3875.2 \pm 0.7_{-1.6}^{+0.3} \pm 0.8 \text{ MeV}/c^2$) and BaBar ($3875.1_{-0.5}^{+0.7} \pm 0.5 \text{ MeV}/c^2$). Both groups also measured branching fractions that are 10 times that for the decay $B \rightarrow KX(3872)$; $X(3872) \rightarrow J/\psi\pi^+\pi^-$. This might be trouble for the $D\bar{D}^*$ mesonic-molecule interpretation [4] of the $X(3872)$, for which the predicted BF ratio is more like 0.1.

Here we report results from a three-times-larger data sample of 657 million $B\bar{B}$ pairs and uses both D^* decays to $D^0\gamma$ and $D^0\pi^0$ to see whether the threshold $D\bar{D}^*$ mass peak is really at a higher mass than $3872 \text{ MeV}/c^2$. We also fit the signal with a more sophisticated form that uses a Breit Wigner and the Flatte fitting function. The Flatte function is essential if the resonance peak mass is below the $D\bar{D}^*$ mass threshold. Fits assuming that the peak is below or above the threshold give the same signal significance of 8.8σ and a fitted peak mass of the Breit Wigner function of $3872.6_{-0.4}^{+0.5} \pm 0.4 \text{ MeV}/c^2$, which is consistent with the current world averaged value for the $X(3872)$ mass from the $J/\psi\pi\pi$ channel. We also measure the product branching fraction for the $D\bar{D}^*$ mode to be $(0.73 \pm 0.17 \pm 0.13) \times 10^{-4}$.

In addition, we searched for the $X(3872)$ in the $J/\psi\pi^+\pi^-$ mass distribution for the $B^0 \rightarrow J/\psi\pi^+\pi^-K^+\pi^-$ decays [6]. Figure 3 a) shows a 5σ significance $X(3872)$ signal in the $J/\psi\pi^+\pi^-$ mass distribution from $B^0 \rightarrow J/\psi\pi^+\pi^-K^+\pi^-$ decays and Fig. 3 b) shows the $K\pi$ mass distribution for the $X(3872)$ mass region. Here the non-resonant $K\pi$ component, shown by the dash-dotted curve, is much stronger than the resonant $K^*(892)^0$ portion shown by the dotted curve. The measured product branching fraction is $B(B^0 \rightarrow X(3872)(K^+\pi^-)_{\text{non-res}}) \times B(X(3872) \rightarrow J/\psi\pi^+\pi^-) = (8.1 \pm 2.0_{-1.4}^{+1.1}) \times 10^{-6}$ at the 90% CL and $B(B^0 \rightarrow X(3872)(K^*(890)^0) \times B(X(3872) \rightarrow J/\psi\pi^+\pi^-) < 3.4 \times 10^{-6}$. This feature is in contrast to corresponding values for conventional charmonium ($c\bar{c}$) states such as $J/\psi, \psi'$ and χ_{c1} , where the $B \rightarrow (c\bar{c})K^*(890)^0$ decay is much larger than non-resonant $B \rightarrow (c\bar{c})K^+\pi^-$ contributions.

We also report the first statistically significant observation of $B^0 \rightarrow X(3872)K_S^0$. The ratio of branching fractions is measured to be $\frac{B^0 \rightarrow X(3872)K_S^0}{B^0 \rightarrow X(3872)K^+} = 0.82 \pm 0.22 \pm 0.05$, which is consistent with unity. The mass difference between the $X(3872)$ produced in B^+ and B^0 decays is measured to be $\Delta M = M_{XK^+} - M_{XK^0} = 0.18 \pm 0.89 \pm 0.26 \text{ MeV}/c^2$, which is consistent with 0.




Figure 4: The $M(\Lambda_c^+ \Lambda_c^-)$ distribution with $\bar{p}$ tag in the $e^+e^- \rightarrow \Lambda_c^+ \Lambda_c^-$ cross section via initial state radiation. The solid curve shows the fit result of the signal. The dashed curve shows that of the threshold function. The combinatorial background parametrization is shown by the dashed-dotted curve.




Figure 5: a) The $e^+e^- \rightarrow \phi\pi^+\pi^-$ cross section distribution fit with two coherent BW functions. b) The $e^+e^- \rightarrow \phi f_0(980)$ cross section distribution fit with one resonance and a coherent nonresonant term.

4. Y states produced via ISR

The Y states are 1^{--} states produced via initial state radiation (ISR). Both the $Y(4260)$ [7] seen in $J\psi\pi^+\pi^-$ masses and the $Y(4325)$ [9] seen in $\psi'\pi^+\pi^-$ masses were discovered in $e^+e^- \rightarrow \gamma_{ISR}Y$ processes by BaBar. They are not the same state.

Belle confirmed both states with the same decay modes as BaBar's also using ISR. However, Belle observed one more peak in the $\pi\pi J\psi$ channel (denoted by $Y(4040)$) with a mass of $M = 4008 \pm 40_{-28}^{+114}$ MeV/ c^2 and the width of $\Gamma = 226 \pm 44 \pm 87$ MeV/ c^2 with a significance of 7.4σ . This state's peak mass is consistent with the $\psi(4040)$ charmonium state in mass, but its measured width is much larger [8]. Belle also observed an additional peak in the $\pi\pi\psi'$ channel (designated as $Y(4660)$) with mass $M = 4664 \pm 11 \pm 5$ MeV/ c^2 & width $\Gamma = 48 \pm 15 \pm 10$ MeV/ c^2 , and a significance of 5.8σ [10]. The $Y(4660)$ also does not fit neatly to any available charmonium assignments. Among these results, one interesting feature is that the $\pi\pi\psi'$ peak masses are different from those seen in the $\pi\pi J/\psi$ channel.

In addition, Belle observed a new near-threshold enhancement in the $e^+e^- \rightarrow \Lambda_c^+ \Lambda_c^-$ cross section using initial-state radiation [11]. Figure 4 shows a clear peak near the $\Lambda_c^+ \Lambda_c^-$ invariant mass threshold. Here only one Λ_c is reconstructed and the other is inferred kinematics when only the decay antiproton is detected. The signal function is a sum of a relativistic S -wave Breit-Wigner function plus a threshold function to account for non-resonant production. The fit result gives a mass and width of $M = 4634_{-7}^{+8}(\text{stat})_{-8}^{+5}(\text{syst})$ MeV/ c^2 and $\Gamma = 92_{-24}^{+40}(\text{stat})_{-21}^{+10}(\text{syst})$ MeV/ c^2 , respectively. These values are consistent with the $Y(4660)$ mass and width. This state can be interpreted as a dibaryon threshold effect such as the one first reported by the BES experiment in the $M(p\bar{p})$ from the decay $J/\psi \rightarrow \gamma p\bar{p}$ [12]. However, the possibility of its being the 5^3S_1 charmonium state is not excluded.

An interesting question is whether or not there exist any XYZ counterparts containing $s\bar{s}$ or $b\bar{b}$ quarks. Here we mention a possible candidate in $s\bar{s}$ sector. The $Y(2175)$ is a 1^{--} state first reported by BaBar in the $f_0\phi$

mass produced in $e^+e^- \rightarrow \gamma_{ISR} f_0 \phi$ ISR events; the mass and width are $M = 2170 \pm 10 \pm 15 \text{ MeV}/c^2$ and $\Gamma = 58 \pm 16 \pm 20 \text{ MeV}/c^2$ [13]. Its production and decay properties are similar as that of $Y(4260)$ and this has stimulated speculation that it may be its $s\bar{s}$ counterpart, the $Y_s(2175)$. This state was confirmed by the BES experiment in $J/\psi \rightarrow \eta \phi f_0(980)$ decays with a mass and width that agree well with the BaBar measurements [14].

We also measure the $\phi\pi^+\pi^-$ and ϕf_0 cross section produced from the $e^+e^- \rightarrow \gamma_{ISR} \phi\pi^+\pi^-$ and $e^+e^- \rightarrow \gamma_{ISR} \phi f_0$ initial state radiation events for the center-of-mass energy ranging from 1.5 to 3.0 GeV [15]. Figure 5 shows corresponding cross section distributions.

A fit with two coherent BW functions is applied to the $\phi\pi^+\pi^-$ cross section distribution; a fit with one BW function interfering with a non-resonant continuum function is applied to the $\phi f_0(980)$ distribution. By averaging the fit results near the mass of the $Y(2175)$, we obtain a mass and width of $2133^{+69}_{-115} \text{ MeV}/c^2$ and $169^{+105}_{-92} \text{ MeV}/c^2$, respectively. The large uncertainty includes both statistical and systematic errors, where the systematic error includes the difference of measured masses from $\phi\pi^+\pi^-$ and ϕf_0 sample as well. This mass agrees with both the BaBar and BES measurements, albeit with a somewhat larger width. The fit to the low-mass peak in Fig. 5 a) gives a mass and width of $M = 1687 \pm 21 \text{ MeV}/c^2$ and $\Gamma = 212 \pm 29 \text{ MeV}/c^2$, respectively, indicating that this is the $\phi(1680)$ radially excited $s\bar{s}$ state. The possibility remains that the $Y(2175)$ could be just a higher radially excited $s\bar{s}$ state.

5. Summary

We observe two more charmonium-like states, Z_1^+ and Z_2^+ , with non-zero charge where both decay into $\chi_{c1}\pi^+$. Their quark content is $c\bar{c}u\bar{d}$. The $X(3872)$ decaying into $\pi^+\pi^-J/\psi$ has been well established, while its underlying nature has not been conclusively identified. The favored interpretations are either a mesonic molecule or a diquark-diantiquark tetraquark meson. Some additional properties of the $X(3872)$ are measured. The mass of the $X(3872)$ decaying into the $D\bar{D}^0$ final state is consistent with that from $\pi^+\pi^-J/\psi$ within errors. Another feature is that the non-resonant $K^+\pi^-$ component in $B^0 \rightarrow X(3872)K^+\pi^-$; $X(3872) \rightarrow J/\psi\pi^+\pi^-$ decay dominates over the resonant $K^*(892)^0$ component. This is not similar to the behaviour usually seen for conventional charmonium states. Among several 1^{--} Y states produced via initial state radiation, we observe the $Y(2175)$ as a possible $s\bar{s}$ analogue to the $Y(4260)$ which, in turn, is a prime candidate for a $c\bar{c}$ -gluon hybrid. The $Y(4660)$ is also seen via its $\Lambda_c^+\Lambda_c^-$ decay mode.

References

- [1] S.-K. Choi, S.L. Olsen et al. (Belle Collaboration), Phys. Rev. Lett. 100, 14200 (2008).
- [2] R. Mizuk, R. Chistov et al. (Belle Collaboration), Phys. Rev. D78, 072004 (2008).
- [3] S.-K. Choi, S.L. Olsen et al. (Belle Collaboration), Phys. Rev. Lett. 91, 262001 (2003).
- [4] E.S. Swanson, Phys. Lett. B588, 189 (2004).
- [5] I. Adachi et al. (Belle Collaboration), arXiv:0810.0358[hep-ex]
- [6] I. Adachi et al. (Belle Collaboration), BELLE-CONF-0849
- [7] B. Aubert et al. (BaBar Collaboration), Phys. Rev. Lett. 95, 142001 (2006).
- [8] C.Z. Yuan et al. (Belle Collaboration), Phys. Rev. Lett. 99, 182004 (2007).
- [9] B. Aubert et al. (BaBar Collaboration), Phys. Rev. Lett. 98, 212001 (2007).
- [10] X.L. Wang et al. (Belle Collaboration), Phys. Rev. Lett. 99, 142002 (2007).
- [11] G. Pakhlova et al. (Belle Collaboration), arXiv:0807.4458[hep-ex], to appear in PRL.
- [12] J.Z. Bai et al. (BES Collaboration), Phys. Rev. Lett. 91, 022001 (2003).
- [13] B. Aubert et al. (BaBar Collaboration), Phys. Rev. D74, 091103 (2006).
- [14] M. Ablikim et al. (BES Collaboration), Phys. Rev. Lett. 100, 102003 (2008).
- [15] I. Adachi et al. (Belle Collaboration), arXiv:0808.0006[hep-ex].